

LINEAR COLLIDER SIGNALS OF ANOMALY MEDIATED SUPERSYMMETRY BREAKING

SOUROV ROY

*Department of Theoretical Physics, Tata Institute of Fundamental Research
Homi Bhabha Road, Mumbai - 400 005, INDIA
E-mail: sourov@theory.tifr.res.in*

*Present address : Department of Physics, Technion, 32000 Haifa, Israel
E-mail: roy@physics.technion.ac.il*

Diagnostic signals of the minimal model of anomaly mediated supersymmetry breaking are discussed in the context of a $\sqrt{s} = 1$ TeV e^+e^- linear collider.

1 Introduction

The idea of anomaly mediated supersymmetry breaking (AMSB) ^{1,2}, where the SUSY breaking is conveyed to the observable sector by the super-Weyl anomaly, has attracted a lot of attention in recent times. This can lead to characteristically distinct and unique signatures in various collider experiments ^{3,4,5}. In this work, we have concentrated on the signals of AMSB which can be studied ^{6,7} in a high energy e^+e^- linear collider at a c.m. energy $\sqrt{s} = 1$ TeV. Anomaly mediation is a special case of gravity mediation with no tree-level couplings between the superfields of the hidden and the observable sectors. This is realized, for instance, when the two sectors are localized on two parallel but distinct 3-branes located in a higher dimensional bulk. This scenario has several distinct and unique features with important phenomenological consequences. The gravitino is rather massive ($m_{3/2} \sim$ tens of TeV), the lightest supersymmetric particle (LSP, $\tilde{\chi}_1^0$) is predominantly a Wino and it is nearly mass-degenerate (mass-splitting $\Delta M < 1$ GeV) with the lighter chargino ($\tilde{\chi}_1^\pm$) which is also a near-Wino. Such a chargino will be long lived and is likely to leave a heavily ionizing charged track in the detector and/or a associated soft pion from the decay $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 + \pi^\pm$. The most glaring problem of AMSB is that it predicts negative mass squares for the sleptons. In the minimal AMSB model this problem is solved by adding a universal constant term m_0^2 to the expressions for squared scalar masses. The set of parameters which define the minimal AMSB model is $\{m_{3/2}, m_0, \text{sign}(\mu), \tan\beta\}$. The AMSB sparticle mass spectrum can be broadly classified into two categories:

- Spectrum A: $\tilde{\chi}_1^0 (\approx \tilde{\chi}_1^\pm) < \tilde{\nu} < \tilde{e}_R (\approx \tilde{e}_L) < \tilde{\chi}_2^0$

Table 1. Possible one or multilepton signal with one soft pion.

Spectrum	Signals	Parent Channels
A	$e \pi$	$\tilde{\nu}\tilde{\nu}, \tilde{e}_L\tilde{e}_L, \tilde{e}_L\tilde{e}_R, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$\mu \pi$	$\tilde{\nu}\tilde{\nu}, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$e e \ell \pi$	$\tilde{e}_R\tilde{e}_R, \tilde{e}_L\tilde{e}_R, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$\mu \mu \ell \pi$	$\tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$\ell_1 \ell_1 \ell_2 \ell_2 \ell_3 \pi$	$\tilde{\chi}_2^0\tilde{\chi}_2^0 (\ell_{1,2,3} = e, \mu)$
B	$e \pi$	$\tilde{\nu}\tilde{\nu}, \tilde{e}_L\tilde{e}_L, \tilde{e}_L\tilde{e}_R, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$\mu \pi$	$\tilde{\nu}\tilde{\nu}, \tilde{e}_L\tilde{e}_L, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$e \ell_1 \ell_2 \pi$	$\tilde{e}_R\tilde{e}_R, \tilde{e}_L\tilde{e}_R, \tilde{e}_L\tilde{e}_L, \tilde{\nu}\tilde{\nu}, \tilde{\chi}_2^0\tilde{\chi}_2^0$
	$\mu \mu \mu \pi$	$\tilde{\chi}_2^0\tilde{\chi}_2^0, \tilde{\nu}\tilde{\nu}$
	$e e \ell_1 \ell_1 \ell_2 \pi$	$\tilde{e}_L\tilde{e}_L, \tilde{e}_R\tilde{e}_R, \tilde{e}_L\tilde{e}_R (\ell_{1,2} = e, \mu)$

- Spectrum B: $\tilde{\chi}_1^0 (\approx \tilde{\chi}_1^\pm) < \tilde{\chi}_2^0 < \tilde{\nu} < \tilde{e}_R (\approx \tilde{e}_L)$.

We discuss our signals for them separately.

2 Results

A list of all possible final states for both the spectra mentioned above, is given in Table 1 for one pion channels. The signals are always accompanied by missing energy. A similar list for two pion channels is given in Ref. ⁷. Cross sections for the various production processes have been calculated for two values of $\tan\beta$, namely 10 and 30 and for $\mu > 0$. In most of the allowed parameter space the signal cross sections are ~ 10 -100 fb. For example, in the worst cases of the signal cross sections, assuming an integrated luminosity of 500 fb^{-1} , one would expect 13165 signal events from spectrum A, while 2910 signal events are predicted from spectrum B for the signal $e^\pm + \pi^\mp + \cancel{E}_T$. The kinematic distributions of the final state particles for the same signal have been studied for a sample point in the AMSB parameter space corresponding to Spectrum A. They are shown in Fig. 1. The chargino decay length distribution shows that a substantial number of events do have a large decay length so that the charged track can be seen. The signals analyzed here are essentially free of Standard Model background.

3 Conclusion

We have investigated possible signals of minimal AMSB model in a 1 TeV c.m energy e^+e^- linear collider. The model is characterized by nearly degenerate lightest neutralino ($\tilde{\chi}_1^0$) and the lighter chargino ($\tilde{\chi}_1^\pm$), resulting in a heavily




Figure 1. Normalized kinematic distributions of decay products: (a) p_T of charged lepton, (b) decay length of the lighter chargino, and (c) p_T of the charged pion arising from $e^\pm + \pi^\mp + \cancel{E}_T$ signal for spectrum A. The AMSB input parameters are $m_{3/2} = 44$ TeV, $\tan \beta = 30$, $\mu > 0$ and $m_0 = 410$ GeV.

ionizing charged track and/or a detectable soft $\pi^\pm$ from the long-lived decay $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 + \text{soft } \pi^\pm$. The generated events triggered by fast charged leptons are large in number and the allowed region of the parameter space can be comprehensively probed.

Acknowledgments

I thank the organisers of SUSY'01 at JINR, Dubna, Russia for the hospitality.

References

1. G.F. Giudice, M.A. Luty, H. Murayama and R. Rattazzi, *J. High Energy Phys.* **12**, 027 (1998).
2. L. Randall and R. Sundrum, *Nucl. Phys. B* **557**, 79 (1999).
3. J.L. Feng, T. Moroi, L. Randall, M. Strassler and S. Su, *Phys. Rev. Lett.* **83**, 1731 (1999).

4. T. Gherghetta, G.F. Giudice and J.D. Wells, *Nucl. Phys. B* **559**, 27 (1999).
5. J.L. Feng and T. Moroi, *Phys. Rev. D* **61**, 095004 (2000).
6. D.K. Ghosh, P. Roy and S. Roy, *J. High Energy Phys.* **08**, 031 (2000).
7. D.K. Ghosh, A. Kundu, P. Roy and S. Roy, *Phys. Rev. D* (to be published), hep-ph/0104217.